



Maxforce[®] GRANULAR

FLY BAIT

ACTIVE INGREDIENT:
Imidacloprid, 1-[6-Chloro-3-pyridinyl)methyl]..... 0.50%
-N-methyl-2-imidazolidinimine..... 0.10%
Z-9-Triicosene..... 99.40%
OTHER INGREDIENTS:..... 100.00%
TOTAL:..... 100.00%
EPA Reg. No. 432-1375 EPA Est. No. 7319-NLD-001

STOP - READ THE LABEL BEFORE USE.
KEEP OUT OF REACH OF CHILDREN.
CAUTION

See side panels for complete First Aid Instructions, Precautionary Statements, Directions for Use and Storage and Disposal Instructions.

Net Weight
40 Pounds (18.14 kg)
SKU# 79848692
US79903316E 140114AV1



Maxforce[®] GRANULAR

FLY BAIT


FIRST AID	
If swallowed:	• Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to by a poison control center or doctor. • Do not give anything to eat or drink.
If on skin:	• Take off contaminated clothing. • Flush with immediately with plenty of water for 15 to 20 minutes. • Call a poison control center or doctor for treatment advice.
If inhaled:	• Have person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. • Call a poison control center or doctor for further treatment advice.
If in eyes:	• Hold eyes open and rinse slowly and gently with water for 15 to 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eyes. • Call a poison control center or doctor for treatment advice.

HAZARDS TO HUMANS AND DOMESTIC ANIMALS
CAUTION
Harmful if swallowed, absorbed through skin, or inhaled. Avoid contact with eyes, skin or clothing. Wash thoroughly with soap and water after handling and before eating, drinking or using tobacco.
ENVIRONMENTAL HAZARDS
This product is highly toxic to aquatic invertebrates.
Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of this product. This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are collecting nectar or pollen from these plants. This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

PRODUCT FACTS	
• Fast acting • Ready-to-use scatter bait or bait-in application. • Contains 10 specially formulated mixtures of attractants • Contains flies resistant to organophosphates and carbamates.	
MAXFORCE GRANULAR FLY BAIT	• Ready-to-use bait containing 0.5% imidacloprid. • Contains the attractant muscarine. • Contains Brevi® lifesting agent to help prevent ingestion by animals, children and pets.
CONTROLS	• Free
WHERE TO APPLY	• Around the outside of structures or partially enclosed or protected areas. • In and around agricultural production facilities. • Around the outside of commercial facilities.
MARKING INFORMATION	• Ready-to-use scatter bait. • Bait-in application. Mix the fly bait with water.
HOW TO APPLY	• Scatter application. • Place in any fly bait station. • Paint-on application.
COVERAGE	17,000 Square Feet (5.7 to 6.3 gal/1000 sq ft)
RE-APPLY	Re-apply every 7 days as needed.
QUESTIONS ??	For questions or comments, call toll-free 1-800-331-2867.

DIRECTIONS FOR USE	
It is a violation of Federal law to use this product if it is not consistent with its labeling.	
WHERE TO APPLY	FLY MANAGEMENT • For use only around the outside of commercial facilities like canneries, dairy, meat and poultry processing plants, beverage plants, food processing plants, restaurants, taverns, supermarkets, warehouses, distribution, greenhouses, recycling areas, loading docks. • For use only around the outside of agricultural production facilities like broiler houses, hatching, livestock housing structures, and farm stables. • For use only around the outside of and in walkways of caged layer houses. Place bait in areas inaccessible to animals. • In horse stables, apply bait only in fly bait stations. Place bait stations in areas inaccessible to animals.
USE RATE	• 5.7 to 6.3 gal/1000 sq ft (scatter bait and bait station) • Apply the bait in many locations. Avoid use in windy locations because flies prefer warm spots for resting. • Time applications to begin at the start of the season before fly populations have reached their peak. • Scatter application: Scatter the bait directly from the container onto dry level surfaces so that the individual granules lie near each other without forming small piles. • Bait stations: Place in any fly bait station. Use 1 bait station to cover approximately 250 sq ft, and add up to 1.6 oz of bait per station. Secure bait stations at least 4 feet above the ground. • Paint-on application: Mix 1.5 oz bait with 1.4 oz warm water (3 lbs bait on 1 quart water) and stir thoroughly. Let stand for about 15 minutes until a paste consistency suitable for painting has formed. Apply the paste with a brush to surfaces where flies rest avoiding surfaces that are dirty. • Restrictions: Do not place bait in areas accessible to food producing animals, children and pets. • Do not allow bait to come into contact with food or water. • Do not apply to milking equipment. • Do not apply inside food processing plants, restaurants or other commercial facilities.

Produced for:
Bayer Environmental Science
A Division of Bayer CropScience LP
2 T. W. Alexander Drive
Research Triangle Park, NC 27709
Product of The Netherlands



Bayer



Maxforce[®] GRANULAR


FLY BAIT

Net Weight
40 Pounds (18.14 kg)
SKU# 79848692
US79903316E 140114AV1




Maxforce[®] GRANULAR

FLY BAIT



Net Weight
40 Pounds (18.14 kg)
SKU# 79848692

Produced for:
Bayer Environmental Science
A Division of Bayer CropScience LP
2 T. W. Alexander Drive
Research Triangle Park, NC 27709
Product of The Netherlands





Maxforce[®]

GRANULAR

FLY BAIT

ACTIVE INGREDIENT:

Imidacloprid, 1-[(6-Chloro-3-pyridinyl)methyl]

-N nitro-2-imidazolidinimine 0.50%

Z-9-Tricosene 0.10%

OTHER INGREDIENTS: 99.40%

TOTAL: 100.00%

EPA Reg. No. 432-1375

EPA Est. No. 7319-NLD-001

STOP - READ THE LABEL BEFORE USE.

KEEP OUT OF REACH OF CHILDREN.

CAUTION

See side panels for complete First Aid Instructions, Precautionary Statements, Directions for Use and Storage and Disposal Instructions.

Net Weight

40 Pounds (18.14 kg)

SKU# 79848692

US79903316E 140114AV1

FIRST AID

If swallowed:	<ul style="list-style-type: none">• Call a poison control center or doctor immediately for treatment advice.• Have person sip a glass of water if able to swallow.• Do not induce vomiting unless told to by a poison control center or doctor.• Do not give anything to an unconscious person.
If on skin:	<ul style="list-style-type: none">• Take off contaminated clothing.• Rinse skin immediately with plenty of water for 15 to 20 minutes.• Call a poison control center or doctor for treatment advice.
If inhaled:	<ul style="list-style-type: none">• Move person to fresh air.• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible.• Call a poison control center or doctor for further treatment advice.
If in eyes:	<ul style="list-style-type: none">• Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.• Call a poison control center or doctor for treatment advice.
Have the product container or label with you when calling a poison control center or doctor or going for treatment. The Bayer Environmental Science Emergency Response Telephone No. is 1-800-334-7577.	
NOTE TO PHYSICIAN: No specific antidote is available. Treat the patient symptomatically.	

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Harmful if swallowed, absorbed through skin, or inhaled. Avoid contact with eyes, skin or clothing. Wash thoroughly with soap and water after handling and before eating, drinking or using tobacco.


ENVIRONMENTAL HAZARDS

This product is highly toxic to aquatic invertebrates.

Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters or rinsate. This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area. This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

PRODUCT FACTS

- Fast acting • Ready-to-use scatter bait or paint-on application
- Contains a specially formulated mixture of attractants • Controls flies resistant to organophosphates and carbamates

MAXFORCE GRANULAR FLY BAIT	<ul style="list-style-type: none">• Ready-to-use bait containing 0.5% imidacloprid.• Contains the attractant muscalure.• Contains Bitrex® bittering agent to help prevent ingestion by animals, children and pets.
CONTROLS	Flies
WHERE TO APPLY	<ul style="list-style-type: none">• Around the outside of structures or partially enclosed or protected areas.• In and around agricultural production facilities.• Around the outside of commercial facilities.
MIXING INFORMATION	<ul style="list-style-type: none">• Ready-to-use scatter bait.• Paint-on application: Mix the fly bait with water.
HOW TO APPLY	<ul style="list-style-type: none">• Scatter application.• Place in any fly bait station.• Paint-on application.
COVERAGE	112,000 Square feet (5.7 to 6.3 oz/1000 sq ft)
RE-APPLY	Re-apply every 7 days as needed.
QUESTIONS ?? 	For questions or comments, call toll-free 1-800-331-2867.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

FLY MANAGEMENT

WHERE TO APPLY

- For use only around the outside of commercial facilities like canneries, dairy, meat and poultry processing plants, beverage plants, food processing plants, restaurants, bakeries, supermarkets, warehouses, dumpsters, grease pits, recycling areas, loading docks.
- For use only around the outside of agricultural production facilities like broiler houses, feedlots, livestock housing structures, and horse stables.
- For use only around the outside of and in walkways of caged layer houses. Place bait in areas inaccessible to animals.
- In horse stables, apply bait only in fly bait stations. Place bait stations in areas inaccessible to animals.

USE RATE

- 5.7 to 6.3 oz/1000 sq ft (scatter bait and bait station)

HOW TO APPLY

Apply the bait in many locations. Avoid cool or windy locations because flies prefer warm sites for resting.

Time applications to begin at the start of the season before fly populations have reached their peak.

- **Scatter application:** Scatter the bait directly from the container onto dry level surfaces so that the individual granules lie near each other without forming small piles.
- **Bait stations:** Place in any fly bait station. Use 1 bait station to cover approximately 250 sq ft, and add up to 1.6 oz of bait per station. Secure bait stations at least 4 feet above the ground.

Bait stations should be inaccessible to food-producing animals, children and pets.

- **Paint-on application:** Mix 1.5 oz bait with 1 fl oz warm water (3 lbs bait in 1 quart water) and stir thoroughly. Let stand for about 15 minutes until a paste consistency suitable for painting has formed. Apply the paste with a brush to surfaces where flies rest avoiding surfaces that are dusty.

RESTRICTIONS



- Do not place bait in areas accessible to food producing animals, children and pets.
- Do not allow bait to come into contact with food or water.
- Do not apply to milking equipment.
- Do not apply inside food processing plants, restaurants or other commercial facilities.

Net Weight

40 Pounds (18.14 kg)

SKU# 79848692

MANAGEMENT TIPS

For Best Results: use the bait as part of an overall Integrated Pest Management program.

- Use proper sanitation and manure management.
- Use along with dilutable sprays like TEMPO®.
- Use proper light management.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

Storage

Store in a cool, dry place and in such a manner as to prevent cross contamination with other pesticides, fertilizers, food, or feed. Store in original containers and out of the reach of children, preferably in a locked storage area. Handle and open container in a manner as to prevent spillage.

Pesticide Disposal

Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Handling

Non-refillable container. Do not reuse or refill this container. Completely empty liner by shaking and tapping sides and bottom to loosen clinging particles. Empty residue into application equipment. Then dispose of liner in a sanitary landfill or by incineration if allowed by State and local authorities. If drum is contaminated and cannot be reused, dispose of in the same manner.

Spills

If the container is leaking or material is spilled, carefully sweep up spilled bait and wash exposed area. In spill or leak incidents, keep unauthorized people away. You may contact the Bayer Environmental Science Emergency Response Team for decontamination procedures or any other assistance that may be necessary. The Bayer Environmental Science Emergency Response Telephone No. is 1-800-334-7577.

IMPORTANT: READ BEFORE USE

Read the entire Directions for Use, Conditions, Disclaimer of Warranties and Limitations of Liability before using this product. If terms are not acceptable, return the unopened product container at once.


By using this product, user or buyer accepts the following Conditions, Disclaimer of Warranties and Limitations of Liability.

CONDITIONS: The directions for use of this product are believed to be adequate and must be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Bayer CropScience LP. All such risks shall be assumed by the user or buyer.

DISCLAIMER OF WARRANTIES: TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BAYER CROPSCIENCE LP MAKES NO OTHER WARRANTIES, EXPRESS OR IMPLIED, OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE OR OTHERWISE, THAT EXTEND BEYOND THE STATEMENTS MADE ON THIS LABEL. No agent of Bayer CropScience LP is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BAYER CROPSCIENCE LP DISCLAIMS ANY LIABILITY WHATSOEVER FOR SPECIAL, INCIDENTAL OR CONSEQUENTIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.

LIMITATIONS OF LIABILITY: TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER IN CONTRACT, WARRANTY, TORT, NEGLIGENCE, STRICT LIABILITY OR OTHERWISE, SHALL NOT EXCEED THE PURCHASE PRICE PAID, OR AT BAYER CROPSCIENCE LP'S ELECTION, THE REPLACEMENT OF PRODUCT.



Bayer (reg'd), the Bayer Cross (reg'd) and MAXFORCE® are registered trademarks of Bayer. Bitrex® is a registered trademark of Johnson Matthey Public Limited Company.



Produced for:
Bayer Environmental Science
A Division of Bayer CropScience LP
2 T. W. Alexander Drive
Research Triangle Park, NC 27709
Product of The Netherlands

Bayer

UN 3077
ENVIRONMENTALLY HAZARDOUS
SUBSTANCE, SOLID, N.O.S.
(IMIDACLOPRID MIXTURE)



Net Weight
40 Pounds (18.14 kg)
SKU# 79848692

Produced for:
Bayer Environmental Science
A Division of Bayer CropScience LP
2 T. W. Alexander Drive
Research Triangle Park, NC 27709
Product of The Netherlands