

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS
AND DOMESTIC ANIMALS

CAUTION

If swallowed, this material may reduce the clotting ability of the blood and cause bleeding. Keep away from children, pets, and domestic animals. Do not contaminate feed and food stuffs. Use only in rodent-infested areas where exposure to agricultural crops or commodities will not occur. Keep away from children, pets, and domestic animals. Wash thoroughly with soap and water after handling and before eating or smoking. Clean clothing should be used daily.

ENVIRONMENTAL HAZARDS

This product is toxic to fish, birds, and other wildlife. Baits exposed to soil surfaces may be hazardous to birds and other wildlife. Do not apply directly to lakes, streams, or ponds.

DIRECTIONS FOR USE

It is a violation of State and Federal Laws to use this product inconsistent with its labeling.

FOR DISTRIBUTION AND USE ONLY IN CALIFORNIA

USE RESTRICTIONS

For use against GROUND SQUIRRELS (*Spermophilus beecheyi*, *S. beecheyi*), DEER MICE (*Peromyscus* spp.), HOUSE MICE (*Mus musculus*).

IMPORTANT

Bait must be eaten at several feedings on 5 or more successive days with no period longer than 48 hours between feedings.

Keep this product in a closed container, free from odors which may reduce acceptability.

BAITING GROUND SQUIRRELS: *Spermophilus beecheyi*, *S. beecheyi*.

Scatter a handful of bait (about 10 baits per pound) evenly over 40 to 50 square feet near active burrows or runways. Repeat every other day for 3 to 4 applications. Do not pile bait!

An uninterrupted supply of bait should be available for 6 to 8 days.

For mechanical spreader applications apply at the rate of 10 pounds per acre through infested area, follow with second application in 2 to 3 days.

BAITING DEER MICE: *Peromyscus* spp.

Spread bait evenly by hand, mechanical spreader, or aircraft at the rate of 2 to 6 pounds per acre through the infested area. Sufficient bait should be applied to last a minimum of 4 days (preferably without rain). In reinfestation programs bait application should precede the sowing of conifer seed by 10 days to 2 weeks.

RODENT BAIT

DIPHACINONE

TREATED GRAIN (0.01 %)

Active Ingredients:

Diphacinone
2-diphenylacetyl-1
3-indandione

Inert Ingredients:
Total:

0.01 %
99.99 %
100.00 %

KEEP OUT OF REACH OF CHILDREN
CAUTION

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: CALL A PHYSICIAN OR POISON CONTROL CENTER IMMEDIATELY. Drink 1 or 2 glasses of water and induce vomiting by touching back of throat. Do not induce vomiting or give anything by mouth to an unconscious person.

NOTE TO PHYSICIAN

Intramuscular and oral administration of Vitamin K, combined with blood transfusions is indicated as in the case of hemorrhage caused by overdoses of bis-hydroxycoumarin (Dicumarol).

REGISTRANT:

California Department of Food and Agriculture
1220 'N' Street, Room A-357
Sacramento, CA 94271-0001

CALIF. REGISTRATION NO.

10965-50003ZA

EPA SLN. NO.

CA890022

EPA EST. NO.

11071-CA-01

NET CONTENTS _____ lbs.

BAITING HOUSE MICE, *Mus musculus*

Place tablespoon amounts (1 4 to 1 2 ounce) of bait in bait box or shallow container, preferably in protected feeder stations. Place bait stations at 8 to 12 foot intervals in dry locations such as in concealed places, in corners, or along walls where house mice feed, drink, or frequent.

Inspect stations daily and add bait as needed. Replace moldy or old bait with fresh bait. Dispose of baits in such a manner as to prevent ingestion by children, pets, wildlife, and domestic animals. An uninterrupted supply of bait should be maintained as long as any bait is taken, which may be 2 to 4 weeks. Where continuous source of infestation is present, permanent bait stations should be established and the bait replenished as needed. All poison baits must be picked up or disposed of after rodent activity has stopped.

STORAGE AND DISPOSAL

Store in a locked enclosure (in a properly posted area) in such a manner so as not to present a hazard to persons, animals, food, feed, crops, or property. Do not contaminate water, food, or feed by storage or disposal. Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Completely empty bag into application equipment. Then, dispose of empty bag in a sanitary landfill or by incineration, or (if allowed by State or local authorities) by burning. If burned, stay out of smoke.

Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.